

Chronic Inflammation

Prolonged tissue injury causes persistent or chronic inflammation.

-Chronic Inflammation may develop:

1. Progressive acute inflammation.
2. Primary chronic inflammation.

Primary chronic inflammation produced by agent

- Dust like carbon, silica, asbestosis
- Microorganisms like virus, bacteria, fungus, and also large parasite.
- Hypersensitivity.
- Unknown causes.

Characteristic features of chronic inflammation are:

1. Infiltration by mononuclear cells
“macrophage, lymphocytes and plasma cells”
2. Proliferation of fibroblasts rather than exudates, The risk of scarring and deformity developing is usually considerably greater than in acute inflammation.

Types of inflammatory cells:

1. Polymorphonuclear leucocytes (PMN)


- *Neutrophil polymorphs (microphages).*
- *Eosinophil polymorphs*
- *Basophils*

2. Lymphoid cells:


- *Lymphocytes*
- *Plasma cell:* This cell is derived from lymphocyte and has small round nucleus in which granules of chromatin are regularly spaced around the periphery giving a (cart-wheel) or (clock face).

Inflammatory Cells

Polymorphnuclear leococytes




Lymphoid and Macrophages





Macrophages

- Special forms of Inflammatory Macrophages:
- *Epithelioid Cells*
- *Giant Cells*

These are epitheloid cells they get their name from the fact they have similar shape to squamous epithelial cells. Their nuclei tend to be long .



Giant cells are a committee of epithelioid or macrophage. Seen here are two giant cells in which the nuclei are lined up around the periphery of the cell.



Effect of chronic Inflammation

- **Beneficial effects:**

1. Proliferation of new blood vessels make the emigration of leukocytes is possible.
2. Fibrous tissue forms barrier to bacteria and toxin.

Side effect

- Is usually accompanied by loss paranchymal cells (tissue destruction).

Repair or Healing

Repair:

Means restitution of injured part of tissue to nearly complete replication of its former structure.

Tissue repair either by:
regeneration or organization.

Regeneration

- Means replacement of tissue cells by proliferation of similar kind of cells.
- Somatic cells are divided into three types according to their ability to regeneration:
- Labile cells: those cells under normal condition continue to divide and multiply throughout the life.
- Stable cells: Normally cease replication when growth ceases but it can retain ability for some regeneration after damage.

Permanent cell

- These cells have no ability to regeneration when damage.
- Organization: It is a process of conversion of inflammatory debris into granulation tissue and finally to the fibrous tissue.

Healing of Skin Wounds

- Healing by first intention “Primary Union”
- Occurs:
- Clean wounds
- Without loss tissues
- The edges of the wound are closed by sutures.

Characterized by:

- Only minimal amount of granulation tissue is formed.
- Take 2-3 weeks for healing.

Healing by second intention

“Secondary Union”

- Occurs:
- Open wounds
- Lost of tissue
- Infected closed wound
- Characterized by:
- Formation of large amount of granulation tissue
- Taking longer time to heal.

Phases of wound healing

- Inflammatory Phase
- Proliferative Phase
- Maturation Phase

Inflammatory phase

- This phase begins at the time of injury. It involves:
- Constriction of injured blood vessels and initiation of blood clotting “blood clot serves as a protect scab”
- After a brief period of constriction, these same vessels dilate and capillaries increase their permeability and produce exudate.

Proliferative phase

- Begins within 2-3 days of injury and may last as long as three weeks in wounds healing by primary union this phase involves:
- Epithelial cells at the margin of the wound begin to regenerate and move toward the center of the wound, forming a new surface layer that destroyed by the injury.
- Proliferation of fibroblasts and newly blood vessels are formed to form a specialized type of soft pink granular tissue called granulation tissue.
- Collagen formed by fibroblast.


Maturation Phase

- Maturation of granulation tissue to form scar tissue.

skin wound healing


HEALING BY FIRST INTENTION

24 hours




Inflammatory phase

3 to 7 days



proliferative phase


Weeks




← maturation phase →

minimal amount of granulation tissue


HEALING BY SECOND INTENTION



Inflammatory phase



proliferative phase



large amount of granulation tissue

Maturation Phase

- Maturation of granulation tissue to form scar tissue.

Maturation Phase

- Maturation of granulation tissue to form scar tissue.

Factors Impair Wound Healing

- Systemic Factors:
 1. Nutritional deficiency e.g. protein deficiency, vitamin C deficiency.
 2. Metabolic diseases like diabetes mellitus.
 3. Poor blood supply.
 4. Hormones such as corticosteroids which act as anti-inflammatory.