

Pathophysiology

=Physiology of altered health

It is mainly concerned with the physiological changes and responses that produce signs and symptoms in case of disease.

So studying of pathophysiology is essential to understanding the rational for medical and surgical therapy.

Disease

- **Inherited disease:** Due to abnormality in the DNA of the fertilized ovum and the cell derived from it.
- **Acquired Disease:** Due to effect of some environmental factors.

Pathogenesis

- It means the mechanism of disease development.

each disease has a characteristic natural history, a typical pattern of evolution, effect and duration that is observed.

Manifestation

- the etiologic agent may provoke a number of changes in the biological process, that lead to produce clinical signs.
- Many diseases have a subclinical stage.

The cell

CHAPTER 1



The Cell



The cell is composed of protoplasm

The protoplasm is composed of:


1. Water (70-85%).
2. Proteins (10-20)% which form cell structures , enzymes.
3. Lipiods (2-3)%
4. Small amounts of carbohydratesare used as fuel.
5. Electrolytes major intracellular electrolytes are, K, Mg, PO₄, SO₄, HCO₃ and small quantity of CL, Na, Ca, Fl.

Cell Membrane

A Carbohydrate chains bound to lipids and proteins




Cell Membrane



cell membrane surrounded the cell and prevent content of the cell get out and prevent molecules from environment to get inside the cell. This membrane is semi-permeable.

- **Component of cell membrane:**
- lipids (phospholipids, and cholesterol)
- Proteins
- Carbohydrate




Cytoplasm

- Cytoplasm is a colloidal solution that contains water, electrolytes, suspended proteins, neutral fats, and glycogen molecules.


In the cytoplasm there are organelles.

Organelles: inner organ of the cell.

Ribosomes



Rough Endoplasmic Reticulum




Rough & Smooth Reticulum




Golgi's Apparatus

Function:


It packages substances (which are synthesized in the cell) into secretory granules.



Mitochondria



Microtubules





The microtubules are slender tubular structures composed of globular proteins called tubulin. Microtubules control cell shape and movement.

Function:

1. Development and maintenance of cell form.
2. Participation in intracellular transport mechanisms.
3. Formation of centrioles.

Microtubules and Centrioles




Centriole


A Microtubule




Electron micrograph of microtubules showing above structural features



B Cilium





C Centriole



Centrioles are cylindrical structure composed of highly organized microtubules. In dividing cells they form the mitotic spindle that aids in the separation and movement of chromosomes.

Nucleus

- It is control center of the cell. It is surrounded by nuclear membrane and it contains the individual units which is called genes.









Genes

- Are units of inheritance which are strung along the chromosomes
- Gene control cell activity by determining the type of protein that is being synthesized in the cytoplasm

Chromosome

- Is a double strand helical molecule of DNA containing variable sequences of four nitrogen bases(thymine, Guanine, Cytosine, and Adenine) these bases form a gene.



Cellular Adaptation

Cells have ability to adapt when there is change in environment or when increase work demands.

Cellular adaptation occurs by:

Changing in size.

Changing in cells number

Changing in cells type.

Atrophy

Decrease in the cell size.

Causes

Disuse

Endocrine stimulation

Malnutrition

Decrease in blood supply.



De- enervation

Atrophy:

a). Brain

This is cerebral atrophy in a patient with Alzheimer's disease. The gyri are narrowed and the sulci widened toward to frontal pole.






Hypertrophy

- Increase in cell size

This is cardiac hypertrophy. The number of myocardial fibers never increases, but their size can increase in response to an increased workload, leading to the marked thickening of the left ventricle in this patient with hypertension.



The main complication of Persistent hypertrophy is *cardiac failure*.



Hyperplasia

Increase in the number of cells in the organ or tissue.

Hyperplasia is a controlled process that occurs in response to an appropriate stimulus and ceases once the stimulus has been removed.

Hyperplasia in the skin

Occurs in case of Psoriasis




FIGURE 1-10

Epidermal hyperplasia. (A) Normal epidermis. (B) Epidermal hyperplasia in psoriasis, shown at the same magnification as in A. The epidermis is thickened, owing to an increase in the number of squamous cells.


Metaplasia

- It means conversion from one tissue to another type like simple epithelium converted to stratified epithelium.

Causes:

- Chronic irritation
- Persist inflammation

IV. Metaplasia



Dysplasia

- Deranged cell growth of a specific tissue that results in cells which vary in size shape and appearance.

Causes

- Chronic irritation

Injury & Cell Death

- Injury:

Any abnormal changes in cell induced by causal agent.

Cell injury is → reversible up to certain point.

→ if the stimulus persists or
sever enough from the beginning

The cell reaches to “point of no return”.

There are two effects of cellular injury:

Cell death:

There is irreversible changes occur in the cell and there is no further integrated function occur like respiration

Lesser form of damage:

Reversible changes and also called degeneration

Cell Death

Cell death means number of cells in the certain tissue are dead.

Somatic death means death of individual (all body).


Necrobiosis: Cell death due to end the physiological age of the cell

Necrosis : Death occurs due to exposure to injurious agent.


Apoptosis: Cell death occurs in case of physiological cell death or due to irreversible cell damage.

Apoptosis

death of single cell by shrinkage of cell, then small parts drop off and are engulfed by macrophage.




1 As a cell begins to undergo apoptosis, its chromosomes condense and its cytoplasm shrinks.





(a)


2 Eventually the nucleus becomes fragmented, its DNA is digested at regular intervals ("laddering"), the cytoplasm becomes fragmented, and the cell extends numerous blebs.




3 Ultimately the remnants of the dead cell (apoptotic bodies) are ingested by phagocytic cells.



(b)




(c)





(d)

NORMAL



Phagocyte



NECROSIS

APOPTOSIS

Necrosis

Causes of

- Marked impaired of blood supply.
- Toxin and chemical poisonous.
- Immunological injury.
- Physical agents.
- Infection.

Necrosis is recognized by:

1). Changes in cytoplasmic staining

The cytoplasm becomes more eosinophilic due to increase affinity to acidic dyes

2). Changes in the nucleus:

a). Pyknosis – condensation of chromatin and shrinkage of the nucleus

b). Karyorrhexis - fragmentation of the nucleus

C). Karyolysis - dissolution of the nucleus by the action of deoxyribonuclease .


2) Changes in cytoplasmic staining:

The cytoplasm becomes more •
eosinophilic due to increase affinity
to acidic dyes

Microscopic appearance of the necrotic tissue:

The tissue appears more eosinophilic than the normal.


Cellular details disappear, so •
there is no nucleus (only
fragments of nucleus may be
seen) and the boundary of the
cell disappears





Types of necrosis:

Coagulative necrosis:

the tissue appears paler than normal adjacent tissue but the architecture of the tissue is maintained.





Gaseous necrosis:

This type of necrosis has a soft cheesy like center.

It most commonly associated with •
T.B.

In this type of necrosis the architecture of the tissue disappears.



Liquefactive necrosis:


Occurs in soft tissues which contains which contain high amounts of fat like brain tissue, and also in case of abscess when large amount or numbers of neutrophils enter the infected area and when neutrophils die that allows lysosomal enzymes to release and digest the tissue, so we can see cavity containing fluids.



The contents of pus are:

Pyogenic bacteria (dead
and life)

Neutrophils (dead and
life)

dead tissue. •





Fat Necrosis

It occurs due to damage to the pancreas which •
leads to release the lipase enzyme. This
enzyme attack fat tissues and lyses fat.

Gangrene

Gangrene

The term gangrene means digestion of dead tissue by saprophytic bacteria (i.e. bacteria which are incapable of invading and multiplying in living tissue). And it associated with foul odor and the color of the tissue changes into dark brown or greenish brown.

Gangrene may be either :

Primary : which necrosis (death) of tissue is due to production of exotoxins by bacteria (which may then invade and digest the dead tissue).

Secondary : which necrosis of tissue is due to other cause like obstruction of blood supply lead to necrosis and then invasion by bacteris.

Primary Gangrene


Like gas gangrene caused by group of bacteria called *Clostridia* especially *Clostridia welchii*, *Clostridium perfringens*, *Clostridia oedematous* and *Clostridia septicum*.

These organisms are intestinal commensals in man and animals.

These organisms are anaerobic and saprophytic.

Their spores are widespread, and are liable to contaminate wounds, but they flourish blood soaked foreign material and dead tissue in dirty puncture or lacerated wounds.

The *Clostridia* produce exotoxins which diffuse into and kill the adjacent tissues and these in turn are invaded



Clostridia produce H₂ and CO₂ which collect as bubbles in the dead tissues, rendering them crepitant on palpation.

Gas gangrene is accompanied by acute hemolysis and sever toxemia.

Secondary Gangrene

Usually result of ischemic necrosis followed by invasion and digestion of dead tissue by putrefactive micro-organisms.

It is most often occur in the foot and leg and in the intestine.

It is of two forms wet and dry gangrene.

Dry gangrene: Occurs in the part of body when there is no excessive fluids like in leg when infraction preceded by gradual arterial occlusion. The skin becomes cold and waxy.

Wet gangrene: Occurs in the part of body when there is excessive fluid ischemic in ischemic area like intestine and edematous leg.

Dry Gangrene





Wet Gangrene



